arXiv:hep-ph/0408133v1 10 Aug 2004

Obtaining α from $B \to \rho^{\pm}\pi^{\mp *}$

J. Zupan [a,b]

[a]Department of Physics, Technion–Israel Institute of Technology, Technion City, 32000 Haifa, Israel

[b]J. Stefan Institute, Jamova 39, P.O. Box 3000,1001 Ljubljana, Slovenia

We discuss how to extract the weak phase α from present data on $B \to \rho^{\pm}\pi^{\mp}$ decays. Introducing $\alpha_{\rm eff}$ and constraining the difference from α using flavor SU(3), one arrives at $\alpha = (95 \pm 16)^{\circ}$, if a testable assumption of a small relative strong phase between the two relevant tree amplitudes is employed to distinguish between discrete ambiguities. On the long run we advocate the combined fit to $B \to \rho^{\pm}\pi^{\mp}$ and SU(3) related modes. The effect of SU(3) breaking in this approach is expected to be small, because of relatively small penguin pollution.

1. Introduction

The difficulty in extracting CKM angle α from time dependent measurements of $b \to u\bar{u}d$ decays is that the penguin contributions to the decays are in general nonnegligible and have to be taken into account. There are two basic approaches to this problem in the case of $\rho^{\pm}\pi^{\mp}$ final state, both dating more than a decade ago. The isospin analysis approach [1,2] is an extension of the original isospin triangle relations approach to extracting α in $B \to \pi\pi$ system [3]. A complication with the $\rho\pi$ final state is that the analysis requires a construction of two pentagons. Since none of the sides of pentagons seem to be much smaller than the others, thus simplifying the analysis, a useful measurement of α using the full isospin analysis may be impractical even with super-B-factory-like luminosities [4].

A more promising way of learning α in these decays is based on performing a time-dependent Dalitz plot analysis of $B^0 \to \pi^+\pi^-\pi^0$ [5], which allows one to obtain the phase differences of decay amplitudes from the interference between two ρ resonance bands. This raises issues such as the precise shapes of the tails of the Breit-Wigner functions, and the effect of interference with other resonant and non-resonant contributions [6]. A complete implementation of this method requires higher statistics than is available today.

In this talk we will try to answer a more modest question, namely what one can learn about α from the data available at this moment? The first obstacle that one encounters in extracting α from the time-dependent decay measurements of $B^0(\overline{B}^0) \to \rho^{\pm}\pi^{\mp}$ by the BABAR [7] and BELLE [8,9] collaborations is that these processes involve more hadronic parameters than measurable quantities. Further assumptions are therefore required to answer the question in a model-independent manner. We will use flavor SU(3), a symmetry less precise than isospin, to relate $B^0 \to \rho^{\pm}\pi^{\mp}$ to processes of the type $B \to K^*\pi$ and $B \to \rho K$ [10,11,12].

2. Observables

We start by setting up notations and conventions. The decay amplitudes are denoted by

$$\begin{aligned} A_{\pm} = A(B^0 \to \rho^{\pm}\pi^{\mp}) &= e^{i\gamma} t_{\pm} + p_{\pm}, \\ \overline{A}_{\pm} = A(\overline{B}^0 \to \rho^{\mp}\pi^{\pm}) &= e^{-i\gamma} t_{\pm} + p_{\pm}. \end{aligned} \tag{1}$$

Each of the four amplitudes can be decomposed in two terms, a "tree" ($t_{\pm}$) and a "penguin" ($p_{\pm}$) amplitude, carrying specific CKM factors. The tree amplitude is proportional to $V_{ub}^* V_{ud}$ by definition, giving dependence on the weak phase γ in Eq. (1). Note that the amplitudes $t_+(p_+)$ and $t_-(p_-)$ have different dynamical origins and are expected to

---

[*] Talk presented at the Sixth International Conference on Hyperons, Charm and Beauty Hadrons, IIT, Chicago, June 27 - July 3, 2004.



Figure 1. The tree (left) and penguin (right) diagrams for the $B^0 \to \rho^+\pi^-$ ($B^0 \to \rho^-\pi^+$) decays.

involve different magnitudes and different strong phases. We define three strong phase differences

$$\delta_\pm = \arg\left(p_\pm/t_\pm\right)\ , \qquad \delta_t = \arg\left(t_-/t_+\right)\ . \quad (2)$$

In addition, we also define the ratios

$$r_\pm \equiv |p_\pm/t_\pm|\ , \qquad r_t \equiv |t_-/t_+|\ . \qquad (3)$$

Counting parameters, we find a total of 8, consisting of 7 hadronic quantities $|t_\pm|, |p_\pm|, \delta_\pm, \delta_t$ and the weak phase γ or α (β is assumed to be known).

Let us now consider measurables in time-dependent rates. Time-dependent decay rates for initially B^0 decaying into $\rho^\pm\pi^\mp$ are given by

$$\Gamma(B^0(t) \to \rho^\pm\pi^\mp) = e^{-\Gamma t}\frac{1}{2}\Gamma^{\rho\pi}\left(1 \pm \mathcal{A}_{\rm CP}^{\rho\pi}\right) \times$$
$$\times\left[1 + (C \pm \Delta C)\cos\Delta m t - (S \pm \Delta S)\sin\Delta m t\right].$$

For initially $\overline{B}^0$ decays, the $\cos\Delta m t$ and $\sin\Delta m t$ terms reverse signs. There are 6 measurables: C, ΔC, S, ΔS, $\Gamma^{\rho\pi}$, $\mathcal{A}_{\rm CP}^{\rho\pi}$ that are parametrized by 8 unknowns discussed above. Thus it is not possible to extract α without further assumptions, which will be made explicit shortly. A compilation of experimental data can be found in [10].

2.1. More economical set

We now introduce a smaller set of observables that still contains all the information needed to extract α. Expanding in terms of penguin-to-tree ratios, $r_\pm$, one arrives at

$$S = \frac{2r_t}{1+r_t^2}\sin 2\alpha\cos\delta_t + O(r_\pm), \qquad (4)$$

$$\Delta S = \frac{2r_t}{1+r_t^2}\cos 2\alpha\sin\delta_t + O(r_\pm). \qquad (5)$$

One can get rid of $|t_\pm|$ (and r_t), if one considers only the ratios of decay widths and appropriately rescaled S, ΔS such that no dependence on r_t occurs. This leads us to the smaller set of 2 direct CP asymmetries

$$\mathcal{A}_{\rm CP}^\pm \equiv \left(|\overline{A}_\pm|^2 - |A_\pm|^2\right) / \left(|\overline{A}_\pm|^2 + |A_\pm|^2\right),$$

and 2 observables related to S and ΔS

$$\begin{Bmatrix}\overline{S}\\ \Delta\overline{S}\end{Bmatrix} \propto \Big[(S+\Delta S)\left(\frac{1+\mathcal{A}_{CP}^{\rho\pi}}{1-\mathcal{A}_{CP}^{\rho\pi}}\right)^{1/2} \pm$$
$$(S-\Delta S)\left(\frac{1-\mathcal{A}_{CP}^{\rho\pi}}{1+\mathcal{A}_{CP}^{\rho\pi}}\right)^{1/2}\Big],$$

where the undisplayed common normalization factor can be found in [10]. Expanding in $r_\pm$ the leading terms are

$$\overline{S} = \sin 2\alpha\cos\delta_t + O(r_\pm), \qquad (6)$$

$$\Delta\overline{S} = \cos 2\alpha\sin\delta_t + O(r_\pm), \qquad (7)$$

$$\mathcal{A}_{\rm CP}^\pm = -\,2r_\pm\sin\delta_\pm\sin(\gamma) + O(r_\pm^2). \qquad (8)$$

The above set of 4 observables depends on 6 unknowns: $\delta_\pm$, $r_\pm$, δ_t, and α. To measure α one has to know $O(r_\pm)$ terms in $\overline{S}$, $\Delta\overline{S}$. There are three basic approaches to this problem, one can either (i) bound corrections, which we will do by constructing $\alpha_{\rm eff}$, (ii) measure corrections, which will be done through SU(3) related modes, or (iii) calculate corrections, for instance in the framework of QCD factorization [13].

3. SU(3) related modes

When relating the tree amplitudes we also account for the SU(3) breaking using guidance from factorization

$$t'_+ = \frac{f_{K^*}}{f_\rho}\frac{V_{ub}^*V_{us}}{V_{ub}^*V_{ud}}\,t_+ = \frac{f_{K^*}}{f_\rho}\bar{\lambda}\,t_+, \qquad (9)$$

$$t'_- = \frac{f_K}{f_\pi}\frac{V_{ub}^*V_{us}}{V_{ub}^*V_{ud}}\,t_- = \frac{f_K}{f_\pi}\bar{\lambda}\,t_-, \qquad (10)$$

while for the penguin amplitudes we assume exact SU(3) relations for the moment

$$p'_\pm = \frac{V_{cb}^*V_{cs}}{V_{cb}^*V_{cd}}\,p_\pm = -\bar{\lambda}^{-1}p_\pm. \qquad (11)$$

Note that in deriving the above relations annihilation like topologies were neglected. Since the tree amplitudes in the $\Delta S = 1$ modes are $\bar{\lambda}$ suppressed, while the penguins are enhanced, they are perfect probes for the sizes of penguin contributions. Of interest are the following ratios of the CP averaged decay widths: ratios with the "tree +penguin" $\Delta S = 1$ decays $\mathcal{R}^0_+ \equiv \bar{\lambda}^2 \overline{\Gamma}(B^0 \to K^{*+}\pi^-)/\overline{\Gamma}(B^0 \to \rho^+\pi^-)$, $\mathcal{R}^0_- \equiv \bar{\lambda}^2 \overline{\Gamma}(B^0 \to \rho^- K^+)/\overline{\Gamma}(B^0 \to \rho^-\pi^+)$, and ratios with the "only penguin" $\Delta S = 1$ decays $\mathcal{R}^+_+ \equiv \bar{\lambda}^2 \overline{\Gamma}(B^+ \to K^{*0}\pi^+)/\overline{\Gamma}(B^0 \to \rho^+\pi^-)$, $\mathcal{R}^+_- \equiv \bar{\lambda}^2 \overline{\Gamma}(B^+ \to \rho^+ K^0)\overline{\Gamma}(B^0 \to \rho^-\pi^+)$, of which only $\mathcal{R}^+_-$ has not been measured yet.

3.1. Bounds on penguins

These ratios can be used to bound $r_\pm$. For r_+ the strictest bound at present is obtained from the "penguin only" ratio $\mathcal{R}^+_+$ that constrains r_+ to be in the range

$$\sqrt{\mathcal{R}^+_+}/\left(1+\sqrt{\mathcal{R}^+_+}\right) \le r_+ \le \sqrt{\mathcal{R}^+_+}/\left(1-\sqrt{\mathcal{R}^+_+}\right),$$

which gives [10]

$$0.14\ (0.16) \le r_+ \le 0.24\ (0.21), \tag{12}$$

at 90 % CL (values in parantheses are obtained by using central values of $\mathcal{R}^+_+$). Symilarly one obtains for r_- [10]

$$0.12(0.21) \le r_- \le 0.32\ (0.27). \tag{13}$$

Note that slightly lower values were predicted in QCD factorization [13]

$$r_+ = 0.10^{+0.06}_{-0.04} \quad \text{and} \quad r_- = 0.10^{+0.09}_{-0.05}. \tag{14}$$

4. α_{eff} and α

Let us now turn to the initial question of bounding α from presently available data. Following $B(t) \to \pi^+\pi^-$ (see e.g. [14]) we define

$$\alpha^\pm_{\text{eff}} \equiv \frac{1}{2} \arg\left(e^{-2i\beta}\overline{A}_\pm A^*_\pm\right), \tag{15}$$

which are *not* observables. The observables on the contrary mix $\overline{A}_+$ and A_- amplitudes and are

$$\begin{aligned} 2\alpha^\pm_{\text{eff}} \pm \hat{\delta} &\equiv \arg\left(e^{-2i\beta}\overline{A}_\pm A^*_\mp\right) \\ &= \arcsin\left(\frac{S \pm \Delta S}{\sqrt{1-(C \pm \Delta C)^2}}\right), \end{aligned} \tag{16}$$

with $\hat{\delta} = \delta_t + O(r_\pm)$. The average of the above observables we define to be α_{eff}

$$\alpha_{\text{eff}} \equiv \frac{1}{2}\left(\alpha^+_{\text{eff}} + \alpha^-_{\text{eff}}\right) = \alpha + O(r_\pm). \tag{17}$$

To have a handle on α we therefore have to estimate $O(r_\pm)$ corrections in (17) which are bounded by SU(3) related modes. The strongest bound in the case of α^+_{eff} comes from "penguin only" ratio, for which at 90% CL

$$\begin{aligned} |\alpha^+_{\text{eff}} - \alpha| &= \frac{1}{2} \arccos\left(\frac{1 - 2\mathcal{R}^+_+ \sin^2(\beta+\alpha)}{\sqrt{1-\mathcal{A}^{+2}_{CP}}}\right) \\ &\le 7.1^\circ - 11.5^\circ. \end{aligned}$$

The equivalent of Charles bound [15] also exists

$$|\alpha^-_{\text{eff}} - \alpha| \le \frac{1}{2} \arccos\left(\frac{1-2\mathcal{R}^0_-}{\sqrt{1-\mathcal{A}^{-2}_{\text{CP}}}}\right) \le 14.9^\circ.$$

Putting the bounds together we have

$$|\alpha_{\text{eff}} - \alpha| \le 11.0^\circ - 13.2^\circ. \tag{18}$$

This implies that we can get a value for α now! Using a mild assumption, testable from partial Dalitz plot analysis [10], that $|\delta_t| \ll 90^\circ$ (QCD factorization gives $\delta_t = (1 \pm 3)^\circ$ [13]) one can distinguish ambiguous solutions for $\alpha^\pm_{\text{eff}} \pm \hat{\delta}$. Namely the difference

$$(2\alpha^+_{\text{eff}} + \hat{\delta}) - (2\alpha^-_{\text{eff}} - \hat{\delta}) = 2\delta_t + \mathcal{O}(r_\pm), \tag{19}$$

has to be small ($\ll 180^\circ$). This leaves the following viable solutions in the $(0^\circ, 180^\circ)$ range

$$\alpha_{\text{eff}} = \{95^\circ \pm 6^\circ, 175^\circ \pm 6^\circ\} \Rightarrow \alpha = (95 \pm 16)^\circ \tag{20}$$

with the last error incorporating also an estimate of SU(3) breaking effects. Note also, that the extracted value of α is most sensitive to changes in S, and not very sensitive to ΔS.

5. SU(3) fit

Finally we discuss how one can obtain α model independently (thus relaxing the mild assumption of $|\delta_t| \ll 90^\circ$, but still neglecting annihilation).



Figure 2. Confidence level (CL) as a function of α for a generated set of data (for details see [10]). Errors used for χ^2 are the currently measured ones [yellow (light gray) region], those anticipated with ten times statistics [cyan (gray)] and hundred times statistics [purple (dark gray)].

Adding SU(3) related modes we have in addition to observables $\overline{S}$, $\Delta\overline{S}$, $\mathcal{A}^{\pm}_{\rm CP}$ also the ratios

$$\frac{1}{2}\left(\mathcal{R}^0_\pm + \mathcal{R}^+_\pm\right) = r^2_\pm + \dots \tag{21}$$

$$\frac{1}{2}\left(\mathcal{R}^+_\pm - \mathcal{R}^0_\pm\right) = r_\pm \bar{\lambda}^2_\pm \cos(\delta_\pm)\cos(\gamma) + \dots \tag{22}$$

giving a total of 8 measurables, and only 6 unknowns: $\delta_\pm$, $r_\pm$, δ_t, α, and therefore consitute an overconstrained system of equations. In Fig. 2 we show an example where data are generated with input parameters $\delta_t = 170^\circ$, $\alpha_{\rm input} = 100^\circ$ and other parameters in experimentally permited ranges (for details see [10]). One sees that (i) there exist ambiguities in the solutions of Eqs. (6)-(8), (21), (22), but (ii) eventually with enough statistics only one solution remains.

The (dis-)apperance of ambiguities can be followed step by step in the $r_\pm$ expansion. At leading order there is a 16-fold ambiguity (cf. Eqs. (6), (7)). This is resolved by higher order terms in $r_\pm$, meaning that the observables have to be measured with $O(r_\pm) \sim 20\%$ precision or better.

A positive aspect of relatively small $r_\pm$, on the other hand, is that the effect of SU(3) breaking on extracted α will be small, effectivelly of order $r^2_\pm$. A Monte Carlo study incorporating up to 30% SU(3) breaking on penguin amplitudes (with flat distribution for the size of the SU(3) breaking) gives $\sqrt{\langle(\alpha^{\rm out} - \alpha^{\rm in})^2\rangle} \sim 2^\circ$ [10].

In conclusion, time dependent measurements of $B(t) \to \rho^\pm\pi^\mp$ allow for a determination of α already with present data, if a mild assumption on the relative tree phase is used, giving

$$\alpha = (95 \pm 16)^\circ. \tag{23}$$

With higher statistics a method that avoids even this mild assumption can be used. The theory error on extracted value of α is very small due to small penguin polution, i.e. $r_\pm \sim 0.2$.

I would like to thank M. Gronau for fruitful collaboration.

REFERENCES

1. H. J. Lipkin, Y. Nir, H. R. Quinn and A. Snyder, Phys. Rev. D **44**, 1454 (1991).
2. M. Gronau, Phys. Lett. B **265**, 389 (1991).
3. M. Gronau and D. London, Phys. Rev. Lett. **65**, 3381 (1990). M. Gronau, D. Pirjol and T. M. Yan, Phys. Rev. D **60**, 03421 (1999); A. J. Buras and R. Fleischer, Eur. Phys. J. C **11**, 93 (1999).
4. J. Stark, eConf **C0304052**, WG423 (2003), hep-ph/0307032.
5. A. E. Snyder and H. R. Quinn, Phys. Rev. D **48**, 2139 (1993); H. R. Quinn and J. P. Silva, Phys. Rev. D **62**, 054002 (2000).
6. A. Deandrea, R. Gatto, M. Ladisa, G. Nardulli and P. Santorelli, Phys. Rev. D **62**, 036001 (2000); A. Deandrea and A. D. Polosa, Phys. Rev. Lett. **86**, 216 (2001); S. Gardner and U. G. Meissner, Phys. Rev. D **65**, 094004 (2002); J. Tandean and S. Gardner, Phys. Rev. D **66**, 034019 (2002).
7. B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **91**, 201802 (2003).
8. A. J. Schwartz, these proceedings.
9. C. C. Wang [Belle Collaboration], hep-ex/0408003.
10. M. Gronau and J. Zupan, hep-ph/0407002.
11. A. Höcker, M. Laget, S. Laplace and J. von Wimmersperg-Toeller, Orsay Report N. LAL-0317 (unpublished).
12. J. Charles *et al.*, hep-ph/0406184.
13. M. Beneke and M. Neubert, Nucl. Phys. B **675**, 333 (2003).
14. M. Gronau, these proceedings, hep-ph/0407316.
15. J. Charles, Phys. Rev. D **59**, 054007 (1999).